

Final Exam Details:

The final exam will take place in room MPHY-205 on May 5th, 2023 from 3:30pm – 5:30pm. Please arrive at least 15 minutes early to get settled; we would like to begin exactly on time. If you have any questions during the exam, you may ask the proctor monitoring the exam.

The exam will consist of 30 multiple-choice questions. You will mark your answers down on the scantron provided and turn in the scantron at the conclusion of the exam.

Chapters to be covered on exam: 1-16**Items to bring:**

- Pencil and eraser
- Scientific Calculator (up to a Ti-84 is allowed)
- Yourself

Items *not* to bring:

- Scantron (one will be provided on exam)
- Computerized Calculator (example: Ti-nspire with touchpad keyboard)
- A laptop, tablet, or phone (all computer devices should be kept powered off and left in your bag for the duration of the exam)
- Any other prohibited item

Useful homework's:

Review through the numbered weekly assignments, and problems covered in recitation/lecture.

Concepts to review:

- Units
 - Size prefix's (examples: kilo-, centi-, mili-, etc.)
 - Conversions (example: kilometers to meters)
 - Significant figures
- Vectors
 - Addition and subtraction
 - Components
 - Magnitude and direction
- Projectile Motion
 - Velocity (1D and 2D)
 - Acceleration (1D and 2D)
 - Falling objects (examples: ball dropped from a height, throwing a rock in the air)
 - Objects moving in a flat plane (examples: block sliding on ice, moving car)
- Newton's laws
 - Force (examples: gravitational force, tension, push & pull)
 - First Law
 - Second Law
 - Third Law

- Free-body Diagrams
- Circular Motion
 - Centripetal force
 - Angular Velocity
 - Angular Acceleration
- Satellite Motion
 - Gravitational force
 - Period
- Work & Energy
 - Definition
 - Kinetic Energy
 - Potential Energy
 - Conservation of Energy
 - Non-conservative Forces
- Momentum
 - Definition
 - Impulse
 - Conservation of Momentum
- Rotational Mechanics
 - Moments
 - Torque
 - Angular momentum
 - Center of Mass
- Fluid Mechanics
 - Quantities
 - Pressure
 - Density
 - Velocity
 - Bernoulli's principle
- Periodic Motion
 - Simple Harmonic Motion
 - Waves
 - Interference
 - Sound
- Thermodynamics
 - Expansion & Heat
 - Thermal Transfer
 - Ideal Gas Law
 - Laws of Thermodynamics
 - Processes
 - Entropy

Formula sheet:**Chapters 1—5 (Exam 1)**

Constant acceleration equations:

$$\begin{aligned} v_x &= v_{0x} + a_x t & x &= x_0 + v_{0x} t + \frac{1}{2} a_x t^2 \\ v_x^2 &= v_{0x}^2 + 2a_x(x - x_0) & x - x_0 &= \left(\frac{v_{0x} + v_x}{2} \right) t \end{aligned}$$

$$g = 9.80 \text{ m/s}^2 \quad w = mg$$

$$\sum F_x = ma_x \quad \sum F_y = ma_y$$

$$f_x = \mu_x n \quad f_x \leq \mu_x n$$

$$F_{\text{spr}} = -kx$$

$$\text{quadratic formula: The equation } ax^2 + bx + c = 0 \text{ has solutions } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

PHYS 201 Formula Sheet**Chapters 9—11 (Exam 3)**For constant α :

$$\begin{aligned} \omega &= \omega_0 + \alpha t & \omega^2 &= \omega_0^2 + 2\alpha(\theta - \theta_0) \\ \theta - \theta_0 &= \omega_0 t + \frac{1}{2} \alpha t^2 & \theta - \theta_0 &= \left(\frac{\omega + \omega_0}{2} \right) t \end{aligned}$$

$$s = r\theta \quad v = r\omega \quad a_{\text{tan}} = r\alpha \quad a_{\text{rad}} = v^2/r = r\omega^2$$

$$K = \frac{1}{2} I \omega^2 \quad I = m_A r_A^2 + m_B r_B^2 + \dots \quad U = M g y_{\text{cm}}$$

$$K_{\text{total}} = \frac{1}{2} M v_{\text{cm}}^2 + \frac{1}{2} I_{\text{cm}} \omega^2$$

$$\tau = F l \quad \sum \tau = I \alpha \quad \Delta W = \tau \Delta \theta \quad P = \tau \omega \quad L = I \omega$$

$$\sum \tau = \frac{\Delta L}{\Delta t} \quad L = m v l$$

first and second conditions for equilibrium:

$$\sum F_x = 0, \sum F_y = 0 \text{ and } \sum \tau = 0 \text{ (any axis)}$$

$$Y = \frac{F_1 / A}{\Delta l / l_0} \quad B = -\frac{\Delta p}{\Delta V / V_0} \quad S = \frac{F_1 / A}{x / h} = \frac{F_1 / A}{\phi}$$

$$F_x = -kx \quad a_x = -\frac{k}{m} x \quad \omega = 2\pi f \quad f = \frac{1}{T}$$

$$U_{\text{el}} = \frac{1}{2} kx^2 \quad K = \frac{1}{2} m v^2$$

$$x = A \cos \omega t \quad v_x = -\omega A \sin \omega t \quad \omega = \sqrt{\frac{k}{m}} \quad f = \frac{1}{2\pi} \sqrt{\frac{k}{m}} \quad T = 2\pi \sqrt{\frac{m}{k}}$$

$$f = \frac{1}{2\pi} \sqrt{\frac{g}{L}} \quad T = 2\pi \sqrt{\frac{L}{g}}$$

Chapters 6—8 (Exam 2)

$$\begin{aligned} a_{\text{rad}} &= \frac{v^2}{R} & v &= \frac{2\pi R}{T} \\ F_x &= G \frac{m_1 m_2}{r^2} & G &= 6.67 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2 & T &= \frac{2\pi r^{3/2}}{\sqrt{G m_E}} \end{aligned}$$

$$W = F_x s = (F \cos \phi) s \quad W_{\text{total}} = K_f - K_i = \Delta K$$

$$U_{\text{grav}} = mgy \quad K = \frac{1}{2} mv^2 \quad U_{\text{el}} = \frac{1}{2} kx^2$$

$$K_f + U_f = K_i + U_i + W_{\text{other}}$$

$$P_{\text{av}} = \frac{W}{t} \quad P = F v$$

$$\vec{p} = m \vec{v} \quad \Delta \vec{p} = \vec{F}_{\text{av}} (t_f - t_i) = \vec{J}$$

$$x_{\text{cm}} = \frac{m_A x_A + m_B x_B + m_C x_C + \dots}{m_A + m_B + m_C + \dots} \quad y_{\text{cm}} = \frac{m_A y_A + m_B y_B + m_C y_C + \dots}{m_A + m_B + m_C + \dots}$$

PHYS 201 Formula Sheet**Chapters 12—16 (Exam 4)**

$$v = f \lambda \quad v = \sqrt{\frac{F_T}{\mu}} \quad y(x, t) = A \sin \left[2\pi f \left(t - \frac{x}{v} \right) \right] = A \sin \left[2\pi \left(\frac{t}{T} - \frac{x}{\lambda} \right) \right]$$

$$f_n = n \left(\frac{v}{2L} \right), \quad n = 1, 2, 3, \dots \quad f_n = n \left(\frac{v}{4L} \right), \quad n = 1, 3, 5, \dots$$

$$I = \frac{P}{4\pi r^2} \quad \beta = (10 \text{ dB}) \log \left(\frac{I}{I_0} \right) \quad f_{\text{beat}} = f_1 - f_2 \quad f_s = \left(\frac{v + v_L}{v + v_s} \right) f_s$$

$$T_F = \frac{9}{5} T_C + 32^\circ \quad T_C = \frac{5}{9} (T_F - 32^\circ) \quad T_K = T_C + 273.15^\circ \quad 1 \text{ C}^\circ = \frac{9}{5} \text{ F}^\circ$$

$$\Delta L = \alpha L_0 \Delta T \quad \Delta V = V_0 \beta \Delta T \quad \frac{F}{A} = -Y \alpha \Delta T$$

$$Q = mc \Delta T \quad Q = \pm mL$$

$$H = kA \frac{T_H - T_C}{L} \quad H = Ae\sigma T^4 \quad \sigma = 5.67 \times 10^{-8} \text{ W/(m}^2 \cdot \text{K}^4\text{)}$$

$$N = 6.022 \times 10^{23} \text{ molecules/mol} \quad m_{\text{total}} = nM$$

$$pV = nRT \quad \rho = \frac{pM}{RT} \quad R = 8.314 \text{ J/(mol} \cdot \text{K)}$$

$$K_u = \frac{3}{2} nRT \quad K_{\text{av}} = \frac{1}{2} m(v^2)_{\text{av}} = \frac{3}{2} kT \quad pV = NkT \quad k = 1.38 \times 10^{-23} \text{ J/K}$$

$$v_{\text{rms}} = \sqrt{\frac{3kT}{m}} = \sqrt{\frac{3RT}{M}} \quad Q = nC \Delta T$$

$$W = p \Delta V \quad W = nRT \ln \left(\frac{V_2}{V_1} \right) \quad \Delta U = Q - W$$

$$C_p = C_v + R \quad p_1 V_1^{\gamma} = p_2 V_2^{\gamma} \quad T_1 V_1^{\gamma-1} = T_2 V_2^{\gamma-1} \quad \gamma = C_p / C_v$$

$$W = Q = |\mathcal{Q}_H| - |\mathcal{Q}_C| \quad e = \frac{W}{\mathcal{Q}_H} = 1 - \frac{|\mathcal{Q}_C|}{|\mathcal{Q}_H|} \quad K = \frac{|\mathcal{Q}_C|}{|\mathcal{W}|} = \frac{|\mathcal{Q}_C|}{|\mathcal{Q}_H| - |\mathcal{Q}_C|}$$

$$\text{Carnot: } \frac{Q_C}{Q_H} = -\frac{T_C}{T_H} \quad e_{\text{Carnot}} = 1 - \frac{T_C}{T_H}$$

$$\Delta S = \frac{Q}{T}$$

$$\rho = \frac{m}{V} \quad p = \frac{F}{A} \quad p = p_{\text{atm}} + \rho g h$$